Syringe

When you select this tool, one or more medicine bottles will appear at the bottom of the screen. A two-step process is required to give the patient an injection. First, you have to draw the medication into the syringe, and then it must be administered.



Drawing medication: After selecting the syringe, place the stylus over the appropriate medicine bottle and slide it upward, as though drawing the medication into the syringe. You may draw medicine in this fashion until the syringe is full.



Administering medication: Place the stylus at the location where you would like to inject the medicine. As long as you hold the stylus against the screen, the medicine will continue to be injected until the syringe is empty. If you lift the stylus before the syringe is empty, you will be able to administer the medicine to another point with whatever remains inside the syringe.

Bandage



At the end of the operation, the final step will be to bandage the sutured incision. After applying the antibiotic gel to arrest the bleeding, select the bandage that appears in the lower right hex. With the bandage selected, place the stylus on one side of the incision and slide it in the direction of the incision to the other side. Lift the stylus to apply the bandage. Make sure it is not too short or too long, and that it is centered over the incision.

Fibrillation

Some patients may undergo cardiac arrest. During fibrillation, the vitals bar will turn red, and the EKG line will begin to shake erratically. The procedure has to come to a complete stop until the heart either resumes its normal pattern or stops beating. If the patient's heart stops, there are two methods that the doctor may employ to resuscitate them:

The Defibrillator

If a patient undergoes cardiac arrest under normal circumstances, the defibrillator will automatically appear; no tool has to be selected. There are two steps to using the defibrillator:



1. Place the stylus on one of the paddles and slide it toward the center of the patient. The other paddle will automatically follow.

2. Once the paddles are set, the voltage meter will appear. Place the stylus on the Touch Screen. When the meter reaches the appropriate level, (designated by a small green section on the meter) lift the stylus from the screen. If the charge is too weak or too strong, the patient's heart may not respond.

Heart Massage

If the defibrillator is unavailable or cannot be used, you'll have to massage the heart. Watch the outlines of the rings, and when they overlap, press the Touch Screen in the center of the rings with the stylus. You'll have to do this successfully a number of times.